

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07D 237/14, 409/04, C07F 9/6509,</b> <b>C07D 237/18, 237/20, A61K 31/50</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/10332</b> <b>(43) International Publication Date:</b> 4 March 1999 (04.03.99)
<b>(21) International Application Number:</b> PCT/US98/17618 <b>(22) International Filing Date:</b> 20 August 1998 (20.08.98)  <b>(30) Priority Data:</b> 08/918,661                      22 August 1997 (22.08.97)                      US  <b>(71) Applicant:</b> ABBOTT LABORATORIES [US/US]; CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).  <b>(72) Inventor:</b> BLACK, Lawrence, A.; 1173 Tamarack Lane, Libertyville, IL 60048 (US).  <b>(74) Agents:</b> WARD, Michael, J. et al.; Abbott Laboratories, CHAD 0377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).		<b>(81) Designated States:</b> CA, JP, MX, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PROSTAGLANDIN ENDOPEROXIDE H SYNTHASE BIOSYNTHESIS INHIBITORS  <b>(57) Abstract</b>  The present invention describes pyridazinone compounds which are cyclooxygenase (COX) inhibitors, and in particular, are selective inhibitors of cyclooxygenase-2 (COX-2). COX-2 is the inducible isoform associated with inflammation, as opposed to the constitutive isoform, cyclooxygenase-1 (COX-1) which is an important "housekeeping" enzyme in many tissues, including the gastrointestinal (GI) tract and the kidneys. The selectivity of these compounds for COX-2 minimizes the unwanted GI and renal side-effects seen with currently marketed non-steroidal anti-inflammatory drugs (NSAIDs).  <div data-bbox="829 1217 1102 1324" style="text-align: right;"><i>Atty. Docket No. 27880A/USA Serial No. 10/803,145 Karen Seibert Reference 103</i></div>		

**Best Available Copy**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## PROSTAGLANDIN ENDOPEROXIDE H SYNTHASE BIOSYNTHESIS INHIBITORS

### Technical Field

The present invention encompasses novel pyridazinone compounds useful  
5 in the treatment of cyclooxygenase-2 mediated diseases. More particularly, this  
invention concerns a method of inhibiting prostaglandin biosynthesis, particularly  
the induced prostaglandin endoperoxide H synthase (PGHS-2, cyclooxygenase-2,  
COX-2) protein.

### Background of the Invention

10 The prostaglandins are extremely potent substances which produce a wide  
variety of biological effects, often in the nanomolar to picomolar concentration  
range. The discovery of two forms of prostaglandin endoperoxide H synthase,  
isoenzymes PGHS-1 and PGHS-2, that catalyze the oxidation of arachidonic acid  
leading to prostaglandin biosynthesis has resulted in renewed research to  
15 delineate the role of these two isozymes in physiology and pathophysiology.  
These isozymes have been shown to have different gene regulation and represent  
distinctly different prostaglandin biosynthesis pathways. The PGHS-1 pathway is  
expressed constitutively in most cell types. It responds to produce prostaglandins  
that regulate acute events in vascular homeostasis and also has a role in  
20 maintaining normal stomach and renal function. The PGHS-2 pathway involves an  
induction mechanism which has been linked to inflammation, mitogenesis and  
ovulation phenomena.


Prostaglandin inhibitors provide therapy for pain, fever, and inflammation,  
and are useful therapies, for example in the treatment of rheumatoid arthritis and  
25 osteoarthritis. The non-steroidal anti-inflammatory drugs (NSAIDs) such as  
ibuprofen, naproxen and fenamates inhibit both isozymes. Inhibition of the  
constitutive enzyme PGHS-1 results in gastrointestinal side effects including ulcers  
and bleeding and incidence of renal problems with chronic therapy. Inhibitors of  
the induced isozyme PGHS-2 may provide anti-inflammatory activity without the  
30 side effects of PGHS-1 inhibitors.

The problem of side-effects associated with NSAID administration has never  
completely been solved in the past. Enteric coated tablets and co-administration  
with misoprostol, a prostaglandin derivative, have been tried in an attempt to  
minimize stomach toxicity. It would be advantageous to provide compounds which  
35 are selective inhibitors of the induced isozyme PGHS-2.

The present invention discloses novel compounds which are selective inhibitors of PGHS-2.

## Summary of the Invention

The present invention discloses pyridazinone compounds which are selective inhibitors of cyclooxygenase-2 (COX-2). The the compounds of the present invention have the formula I:



I

where

X is selected from the group consisting of O, S, NR<sup>a</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>a</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic, heterocyclic (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy, alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl, haloalkynyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, aryl, arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy, arylcarbonylalkyl, heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy), heterocyclic (oxy), -C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -R<sup>6</sup>-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OH)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(NOR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(NR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NOR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NR<sup>d</sup>R<sup>6</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CHX'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(CHX'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>,

wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, haloalkenyl, haloalkynyl, heterocyclic, and heterocyclic (alkyl);

wherein  $R^6$  is alkylene or alkenylene;

5  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl);


$R^{20}$  is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

10  $R^d$  and  $R^e$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl);

$X'$  is halogen;

$n$  is from 0 to about 10, and  $m$  is 0 to about 5;

15 at least one of  $R^1$ ,  $R^2$  and  $R^3$  is a group, substituted with a substituent having a group  $-X^1-R^9$ , having the formula:



where  $X^1$  is selected from the group consisting of  $-SO_2-$ ,  $-SO(NR^{10})-$ ,  $-PO(OR^{11})-$ , and  $-PO(NR^{12}R^{13})-$ ,

20  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino,  $-NHNH_2$ , alkylamino, dialkylamino, alkoxy, thiol, alkylthiol, and protecting groups,

$X^2$  is selected from the group consisting of hydrogen or halogen;

25  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$ , and  $R^{13}$  are independently selected from the group consisting of hydrogen, alkyl, and cycloalkyl, or  $R^{12}$  and  $R^{13}$  can be taken together, with the nitrogen to which they are attached, to form a heterocyclic ring having from 3 to 6 atoms.

The remaining two of the groups of R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup>, are independently selected from the group consisting of hydrogen, hydroxy, hydroxyalkyl, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, amido, amidoalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, amino, aminocarbonyl, aminocarbonylalkyl, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -C(R<sup>16</sup>)(R<sup>17</sup>)-,

C(O)NR<sup>21</sup>-, -C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl),

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; and

R<sup>21</sup> is selected from the group consisting of hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.


## 20 Detailed Description of the Invention

All patents, patent applications, and literature references cited in the specification are hereby incorporated by reference in their entirety. In the case of inconsistencies, the present disclosure, including definitions, will prevail.

The present invention discloses pyridazinone compounds which are cyclooxygenase (COX) inhibitors and are selective inhibitors of cyclooxygenase-2 (COX-2). COX-2 is the inducible isoform associated with inflammation, as opposed to the constitutive isoform, cyclooxygenase-1 (COX-1) which is an important "housekeeping" enzyme in many tissues, including the gastrointestinal (GI) tract and the kidneys.

In one embodiment, the compounds of the present invention have the formula I:

-5-



I

where

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and  
 5 N-NR<sup>b</sup>RC, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl,  
 cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic,  
 heterocyclic (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently  
 selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and  
 arylalkyl;

10 R is selected from the group consisting of hydrogen, alkyl, alkenyl,  
 alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy,  
 alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl, haloalkynyl,  
 hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, aryl,  
 arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy, arylcarbonylalkyl,  
 15 heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy), heterocyclic (oxy), -C(O)R<sup>5</sup>,  
 -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -R<sup>6</sup>-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OH)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(NOR<sup>d</sup>)R<sup>5</sup>,  
 -(CH<sub>2</sub>)<sub>n</sub>C(NR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NOR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NR<sup>d</sup>R<sup>e</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-R<sup>7</sup>,  
 -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sup>3</sup>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sup>3</sup>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sup>2</sup>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sup>3</sup>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sup>3</sup>)]<sub>m</sub>-  
 (CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sup>2</sup>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CHX')<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sup>3</sup>,  
 20 -(CH<sub>2</sub>)<sub>n</sub>(CHX')<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>,

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl,  
 alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, haloalkenyl,  
 haloalkynyl, heterocyclic, and heterocyclic (alkyl);

wherein R<sup>6</sup> is alkylene or alkenylene;

25 R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of  
 hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl,  
 arylalkyl, heterocyclic, and heterocyclic (alkyl);


R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

R<sup>d</sup> and R<sup>e</sup> are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl);

X' is halogen;

n is from 0 to about 10, and m is 0 to about 5;

at least one of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> is a group, substituted with a substituent having a group -X<sup>1</sup>-R<sup>9</sup>, having the formula:



where X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, -SO(NR<sup>10</sup>)-, -PO(OR<sup>11</sup>)-, and -PO(NR<sup>12</sup>R<sup>13</sup>)-,

R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, -NHNH<sub>2</sub>, alkylamino, dialkylamino, alkoxy, thiol, alkylthiol, and protecting groups,

X<sup>2</sup> is selected from the group consisting of hydrogen or halogen;

R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, and R<sup>13</sup> are independently selected from the group consisting of hydrogen, alkyl, and cycloalkyl, or R<sup>12</sup> and R<sup>13</sup> can be taken together, with the nitrogen to which they are attached, to form a heterocyclic ring having from 3 to 6 atoms.

The remaining two of the groups of R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup>, are independently selected from the group consisting of hydrogen, hydroxy, hydroxyalkyl, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, amido, amidoalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, amino, aminocarbonyl, aminocarbonylalkyl, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro,




and  $-Y-R^{14}$ , wherein Y is selected from the group consisting of,  $-O-$ ,  $-S-$ ,  $-C(R^{16})$  ( $R^{17}$ )-,

$C(O)NR^{21}$ -,  $-C(O)-$ ,  $-C(O)O-$ ,  $-NH-$ ,  $-NC(O)-$ , and  $-NR^{19}$ -.  $R^{14}$  is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cyclo-  
 5 alkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl),

$R^{16}$ ,  $R^{17}$ , and  $R^{19}$  are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; and


$R^{21}$  is selected from the group consisting of hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl),  
 10 or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another embodiment, compounds of the present invention have the formula II:



II

wherein Z is a group having the formula:



where  $X^1$  is selected from the group consisting of  $-SO_2-$ ,  $-$  and  $SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino;  
 20

$X^2$  is selected from the group consisting of hydrogen or halogen;

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic, heterocyclic (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy, alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl, haloalkynyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, aryl, arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy, arylcarbonylalkyl, heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy), heterocyclic (oxy), -C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CHX'<sub>1</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(CHX'<sub>1</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>,

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl),

R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

X' is halogen;


n is from 0 to about 10, m is from 0 to about 5;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of hydrogen, hydroxy, hydroxyalkyl, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -C(R<sup>16</sup>) (R<sup>17</sup>)-, -C(O)-, -C(O)O-, -NH-, and

-NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; ; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In yet another embodiment, compounds of the present invention have the formula **III**:



**III**

wherein X, X<sup>1</sup>, R, R<sup>1</sup>, R<sup>3</sup>, and R<sup>9</sup> are as defined in Formula I; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

15

In a preferred embodiment, compounds of the present invention have the formula **III**, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic, heterocyclic (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl,

25

cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl,  $-(CH_2)_n C(O)R^5$ ,  $-(CH_2)_n C\equiv C-R^7$ ,  $-(CH_2)_n (CX'_2)_m CX'_3$ ,  $-(CH_2)_n (CX'R^6)_m CX'_3$ ,  $-(CH_2)_n (CX'_2)_m R^8$ ,  $-(CH_2)_n (CX'R^6)_m R^8$ , and  $-(CH_2)_n R^{20}$ ;

5                    wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl),  $R^{20}$  is selected from the group  
10                   consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

$X'$  is halogen;

$R^6$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxy, and haloalkyl;  
15                  

$n$  is from 0 to about 10,  $m$  is from 0 to about 5;

$R^1$  and  $R^3$  are independently selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and  $-Y-R^{14}$ , wherein  $Y$  is selected from  
20                   the group consisting of,  $-O-$ ,  $-S-$ ,  $-CH_2-$ ,  $-CHR^{15}-$ ,  $-C(R^{16})(R^{17})-$ ,  $-C(O)-$ ,  $-NH-$ , and  $-NR^{19}-$ .  $R^{14}$  is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

25                    $R^{15}$ ,  $R^{16}$ ,  $R^{17}$ , and  $R^{19}$  are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula III, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

5 X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>RC, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

10 R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

15 wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

20 n is from 0 to about 10;

R<sup>1</sup> and R<sup>3</sup> are independently selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>)(R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

25 R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl,

heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula III, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>RC, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, and -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl); and

n is from 0 to about 10;

R<sup>1</sup> and R<sup>3</sup> are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>)(R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula III, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>RC<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup> where R<sup>20</sup> is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

R<sup>1</sup> and R<sup>3</sup> are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>)(R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula III, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X is selected from the group consisting of O, S,  $\text{NR}^4$ ,  $\text{N-OR}^a$ , and  $\text{N-NR}^b\text{R}^c$ , wherein  $\text{R}^4$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and  $\text{R}^a$ ,  $\text{R}^b$ , and  $\text{R}^c$  are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(\text{CH}_2)_n\text{-R}^{20}$  where  $\text{R}^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

$\text{R}^1$  and  $\text{R}^3$  are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl) ; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula III, wherein  $\text{X}^1$  is selected from the group consisting of  $-\text{SO}_2-$ , and  $-\text{SO}(\text{NR}^{10})-$ , and  $\text{R}^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X is selected from the group consisting of O, S,  $\text{NR}^4$ ,  $\text{N-OR}^a$ , and  $\text{N-NR}^b\text{R}^c$ , wherein  $\text{R}^4$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and  $\text{R}^a$ ,  $\text{R}^b$ , and  $\text{R}^c$  are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(\text{CH}_2)_n\text{-R}^{20}$  where  $\text{R}^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;


n is from 0 to about 10;

$\text{R}^1$  and  $\text{R}^3$  are independently selected from the group consisting of hydrogen, phenyl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine including, but not limited to,



*p*-chlorophenyl, *p*-fluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluoro-phenyl, and the like; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In yet another embodiment, compounds of the present invention have formula IV:



IV

where R, R<sup>1</sup>, X<sup>1</sup>, X<sup>2</sup>, R<sup>9</sup>, and R<sup>3</sup> are as described in Formula I; or a pharmaceutically acceptable salt or ester thereof.

In a preferred embodiment, compounds of the present invention have the formula IV, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CX'<sub>2</sub>)<sub>m</sub>CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(CX'R<sup>6</sup>)<sub>m</sub>CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(CX'<sub>2</sub>)<sub>m</sub>R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CX'R<sup>6</sup>)<sub>m</sub>R<sup>8</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl), R<sup>20</sup> is selected from the group

consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

X' is halogen;

R<sup>6</sup> is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, hydroxy, and haloalkyl;

n is from 0 to about 10, m is from 0 to about 5;

R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>)(R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula IV, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>. and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

n is from 0 to about 10;

R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>)(R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula IV, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, and -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl); and

n is from 0 to about 10;

R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>) (R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In another preferred embodiment, compounds of the present invention have the formula IV, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

R is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup> where R<sup>20</sup> is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -CHR<sup>15</sup>-, -C(R<sup>16</sup>) (R<sup>17</sup>)-, -C(O)-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and

R<sup>15</sup>, R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano.

In another preferred embodiment, compounds of the present invention have the formula IV, wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

- 5         $R$  is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

$n$  is from 0 to about 10;

- 10         $R^1$  is hydrogen and  $R^3$  is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl); or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- In another preferred embodiment, compounds of the present invention have the formula IV, wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

$R$  is selected from hydrogen, haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

- 20         $n$  is from 0 to about 10;

- $R^1$  is hydrogen and  $R^3$  is selected from the group consisting of hydrogen, phenyl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine including, but not limited to, *p*-chlorophenyl, *p*-fluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluoro-phenyl, and the like; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- In another preferred embodiment, compounds of the present invention have the formula IV, wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

R is haloalkyl, R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen and phenyl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

5

In another preferred embodiment, compounds of the present invention have the formula IV, wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

10

R is substituted and unsubstituted aryl and R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen and phenyl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

15

In another preferred embodiment, compounds of the present invention have the formula IV, wherein X' is SO<sub>2</sub>, R<sup>9</sup> is selected from alkyl and amino, R is substituted and unsubstituted aryl and R<sup>1</sup> is hydrogen and R<sup>3</sup> is selected from the group consisting of hydrogen and phenyl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

20

## Preferred Embodiments

Compounds useful in practicing the present invention include, but are not limited to:

25

5-(4-Methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Benzyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Methyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
30 2-Ethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;

- 2-(4-Fluorobenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(n-Butyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(2,2,2-Trifluoroethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
5 pyridazinone;  
2-(4-Fluoro- $\alpha$ -methylbenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
pyridazinone;  
2-(n-Propyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(n-Pentyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
10 2-Cyclohexylmethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Phenacyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Propargyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Cyclohexyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
15 2-(2-Butynyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(Cyclobutylmethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
pyridazinone; and  
2-(3-Methylbuten-2-yl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
pyridazinone; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

20

## **Preparation of Compounds of the Invention**


25 The compounds of the invention may be prepared by a variety of synthetic routes. Representative procedures are outlined in Scheme 1 below.

A general route to the compounds of the invention having Formula III, where the aryl group at the 5-position on the pyridazinone ring is substituted with a sulfonyl group ring ( $R^2$  is aminosulfonyl, or methylsulfonyl, shown as methylsulfonyl) is described in Scheme 1, below. Phenyl acetic acid, optionally substituted, was  
30 condensed with a benzaldehyde to form the corresponding 2-phenyl-*trans*-cinnamic acid. The *trans*-cinnamic acid was converted to the corresponding acid chloride which was then decarbonylated to provide the benzyl-aryl ketone. The ketone was


then alkylated with ethyl bromoacetate in the presence of a strong base such as sodium bis(trimethylsilyl)amide. Treatment of the ketoester product with hydrazine, in an alcohol solvent, such as ethanol, provided dihydropyridazinone. The dihydropyridazinone was dehydrogenated to form the pyridazinone by treatment with

5 Bromine in acetic acid. If desired, the R' group can be added via substitution using an appropriate alkylating agent. In Scheme 1, X is defined as in R<sup>3</sup> and R' is defined as in R.

## SCHEME 1







5

## Definitions of Terms

As used throughout this specification and the appended claims, the following terms have the meanings specified.

- 10 The term "protecting groups includes "carboxy protecting group" and "N-protecting groups". "Carboxy protecting group" as used herein refers to a carboxylic acid protecting ester group employed to block or protect the carboxylic acid functionality while the reactions involving other functional sites of the compound are carried out. Carboxy protecting groups are disclosed in Greene,
- 15 "Protective Groups in Organic Synthesis" pp. 152-186 (1981), which is hereby incorporated herein by reference. In addition, a carboxy protecting group can be used as a prodrug whereby the carboxy protecting group can be readily cleaved *in vivo*, for example by enzymatic hydrolysis, to release the biologically active parent. T. Higuchi and V. Stella provide a thorough discussion of the prodrug concept in
- 20 "Pro-drugs as Novel Delivery Systems", Vol 14 of the A.C.S. Symposium Series, American Chemical Society (1975), which is hereby incorporated herein by

reference. Such carboxy protecting groups are well known to those skilled in the art, having been extensively used in the protection of carboxyl groups in the penicillin and cephalosporin fields, as described in U.S. Pat. No. 3,840,556 and 3,719,667, the disclosures of which are hereby incorporated herein by reference.

5 Examples of esters useful as prodrugs for compounds containing carboxyl groups can be found on pages 14-21 of "Bioreversible Carriers in Drug Design: Theory and Application", edited by E.B. Roche, Pergamon Press, New York (1987), which is hereby incorporated herein by reference. Representative carboxy protecting groups are C<sub>1</sub> to C<sub>8</sub> alkyl (e.g., methyl, ethyl or tertiary butyl and the like); haloalkyl;

10 alkenyl; cycloalkyl and substituted derivatives thereof such as cyclohexyl, cyclopentyl and the like; cycloalkylalkyl and substituted derivatives thereof such as cyclohexylmethyl, cyclopentylmethyl and the like; arylalkyl, for example, phenethyl or benzyl and substituted derivatives thereof such as alkoxybenzyl or nitrobenzyl groups and the like; arylalkenyl, for example, phenylethenyl and the like; aryl and substituted derivatives thereof, for example, 5-indanyl and the like;

15 dialkylaminoalkyl (e.g., dimethylaminoethyl and the like); alkanoyloxyalkyl groups such as acetoxymethyl, butyryloxymethyl, valeryloxymethyl, isobutyryloxymethyl, isovaleryloxymethyl, 1-(propionyloxy)-1-ethyl, 1-(pivaloyloxy)-1-ethyl, 1-methyl-1-(propionyloxy)-1-ethyl, pivaloyloxymethyl, propionyloxymethyl and the like;

20 cycloalkanoyloxyalkyl groups such as cyclopropylcarbonyloxymethyl, cyclobutylcarbonyloxymethyl, cyclopentylcarbonyloxymethyl, cyclohexylcarbonyloxymethyl and the like; aroyloxyalkyl, such as benzoyloxymethyl, benzoyloxyethyl and the like; arylalkylcarbonyloxyalkyl, such as benzylcarbonyloxymethyl, 2-benzylcarbonyloxyethyl and the like;

25 alkoxy-carbonylalkyl, such as methoxycarbonylmethyl, cyclohexyloxy-carbonylmethyl, 1-methoxycarbonyl-1-ethyl, and the like; alkoxy-carbonyloxyalkyl, such as methoxycarbonyloxymethyl, t-butyloxy-carbonyloxymethyl, 1-ethoxycarbonyloxy-1-ethyl,

30 1-cyclohexyloxy-carbonyloxy-1-ethyl and the like; alkoxy-carbonylaminoalkyl, such as t-butyloxy-carbonylamino-methyl and the like; alkylaminocarbonylaminoalkyl, such as methylaminocarbonylamino-methyl and the like; alkanoylaminoalkyl, such as acetylaminomethyl and the like; heterocycliccarbonyloxyalkyl, such as 4-methylpiperazinyloxy-carbonyloxymethyl and the like; dialkylaminocarbonylalkyl, such

as dimethylaminocarbonylmethyl, diethylaminocarbonylmethyl and the like; (5-(loweralkyl)-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-t-butyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like; and (5-phenyl-2-oxo-1,3-dioxolen-4-yl)alkyl, such as (5-phenyl-2-oxo-1,3-dioxolen-4-yl)methyl and the like.

- 5       The term "N-protecting group" or "N-protected" as used herein refers to those groups intended to protect the N-terminus of an amino acid or peptide or to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups In Organic Synthesis," (John Wiley & Sons, New York (1981)), which is hereby
- 10       incorporated by reference. N-protecting groups comprise acyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxycarbonyl,  $\alpha$ -chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; carbamate forming
- 15       groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-
- 20       dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl,  $\alpha,\alpha$ -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butyloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-
- 25       methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like; alkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like; and silyl groups such as trimethylsilyl and the like. Preferred N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, phenylsulfonyl, benzyl, t-butyloxycarbonyl (t-Boc)
- 30       and benzyloxycarbonyl (Cbz).

The term "alkanoyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a carbonyl (-C(O)-) group. Examples of alkanoyl include acetyl, propionyl and the like.

5 The term "alkanoylamino" as used herein refers to an alkanoyl group as previously defined appended to an amino group. Examples alkanoylamino include acetamido, propionylamido and the like.

10 The term "alkenyl" as used herein refers to a straight or branched chain hydrocarbon radical containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon double bond. Alkenyl groups include, for example, vinyl (ethenyl), allyl (propenyl), butenyl, 1-methyl-2-buten-1-yl and the like.

The term "alkenylene" denotes a divalent group derived from a straight or branched chain hydrocarbon containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon double bond. Examples of alkenylene include -CH=CH-, -CH<sub>2</sub>CH=CH-, -C(CH<sub>3</sub>)=CH-, -CH<sub>2</sub>CH=CHCH<sub>2</sub>-,  
15 -CH<sub>2</sub>CH=CHCH<sub>2</sub>CH=CHCH<sub>2</sub>-, and the like.

The term "alkenyloxy" as used herein refers to an alkenyl group, as previously defined, connected to the parent molecular moiety through an oxygen (-O-) linkage. Examples of alkenyloxy include allyloxy, butenyloxy and the like.

20 The term "alkoxy" as used herein refers to R<sub>41</sub>O- wherein R<sub>41</sub> is a loweralkyl group, as defined herein. Examples of alkoxy include, but are not limited to, ethoxy, tert-butoxy, and the like.

The term "alkoxyalkoxy" as used herein refers to R<sub>80</sub>O-R<sub>81</sub>O- wherein R<sub>80</sub> is loweralkyl as defined above and R<sub>81</sub> is alkylene. Representative examples of alkoxyalkoxy groups include methoxymethoxy, ethoxymethoxy, t-butoxymethoxy  
25 and the like.

The term "alkoxycarbonyl" as used herein refers to an alkoxyl group as previously defined appended to the parent molecular moiety through a carbonyl group. Examples of alkoxycarbonyl include methoxycarbonyl, ethoxycarbonyl, isopropoxycarbonyl and the like.

The term "alkoxyalkoxyalkenyl" as used herein refers to an alkoxyalkoxy group as previously defined appended to an alkenyl radical. Representative examples of alkoxyalkoxyalkenyl groups include methoxyethoxyethenyl, methoxymethoxymethenyl, and the like.

- 5       The term "alkoxyalkyl" as used herein refers to an alkoxy group as previously defined appended to an alkylene as previously defined. Examples of alkoxyalkyl include, but are not limited to, methoxymethyl, methoxyethyl, isopropoxymethyl and the like.

- 10       The term "(alkoxycarbonyl)thioalkoxy" as used herein refers to an alkoxycarbonyl group as previously defined appended to a thioalkoxy radical. Examples of (alkoxycarbonyl)thioalkoxy include methoxycarbonylthiomethoxy, ethoxycarbonylthiomethoxy and the like.

- 15       The terms "alkyl" and "loweralkyl" as used herein refer to straight or branched chain alkyl radicals containing from 1 to 15 carbon atoms including, but not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, n-pentyl, 1-methylbutyl, 2,2-dimethylbutyl, 2-methylpentyl, 2,2-dimethylpropyl, n-hexyl and the like.

The term "alkylamino" as used herein refers to  $R_{51}NH-$  wherein  $R_{51}$  is a loweralkyl group, for example, ethylamino, butylamino, and the like.

- 20       The term "alkylcarbonylalkyl" as used herein refers to  $R_{40}-C(O)-R_{41}$  wherein  $R_{40}$  is an alkyl group and  $R_{41}$  is alkylene.

- 25       The term "alkylene" denotes a divalent group derived from a straight or branched chain saturated hydrocarbon having from 1 to 15 carbon atoms by the removal of two hydrogen atoms, for example  $-CH_2-$ ,  $-CH_2CH_2-$ ,  $-CH(CH_3)-$ ,  $-CH_2CH_2CH_2-$ ,  $-CH_2C(CH_3)_2CH_2-$  and the like.

The term "alkylsulfonyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a sulfonyl ( $-S(O)_2-$ ) group. Examples of alkylsulfonyl include methylsulfonyl, ethylsulfonyl, isopropylsulfonyl and the like.

The term "alkylsulfonylalkyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a sulfonylalkyl ( $-\text{S}(\text{O})_2-\text{R}_{52}-$ ) group wherein  $\text{R}_{52}$  is alkylene. Examples of alkylsulfonylalkyl include methylsulfonylmethyl ( $\text{CH}_3-\text{S}(\text{O})_2-\text{CH}_2-$ ),

5 ethylsulfonylmethyl, isopropylsulfonylethyl and the like.

The term "alkylsulfonylamino" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a sulfonylamino ( $-\text{S}(\text{O})_2-\text{NH}-$ ) group. Examples of alkylsulfonylamino include methylsulfonylamino, ethylsulfonylamino, isopropylsulfonylamino and the like.

10 The term "alkylsulfonylarylalkyl" as used herein refers to an alkyl group as previously defined appended to the parent molecular moiety through a sulfonylalkyl ( $-\text{S}(\text{O})_2-\text{R}_{45}\text{R}_{33}-$ ) group wherein  $\text{R}_{45}$  is aryl and  $\text{R}_{33}$  is alkylene. Examples of alkylsulfonylarylalkyl include methylsulfonylphenylmethyl ethylsulfonylphenylmethyl, isopropylsulfonylphenylethyl and the like.

15 The term "alkylthio" as used herein refers to  $\text{R}_{53}\text{S}-$  wherein  $\text{R}_{53}$  is alkyl.

The term "alkynyl" as used herein refers to a straight or branched chain hydrocarbon radical containing from 2 to 15 carbon atoms and also containing at least one carbon-carbon triple bond. Examples of alkynyl include  $-\text{C}\equiv\text{C}-\text{H}$ ,  $\text{H}-\text{C}\equiv\text{C}-\text{CH}_2-$ ,  $\text{H}-\text{C}\equiv\text{C}-\text{CH}(\text{CH}_3)-$  and the like.

20 The term "amido" as used herein refers to  $\text{R}_{54}-\text{C}(\text{O})-\text{NH}-$  wherein  $\text{R}_{54}$  is an alkyl group.

The term "amidoalkyl" as used herein refers to  $\text{R}_{34}-\text{C}(\text{O})-\text{NHR}_{35}-$  wherein  $\text{R}_{34}$  is alkyl and  $\text{R}_{35}$  is alkylene.

The term "amino" as used herein refers to  $-\text{NH}_2$ .

25 The term "aminocarbonyl" as used herein refers to  $\text{H}_2\text{N}-\text{C}(\text{O})-$ .

The term "aminocarbonylalkyl" as used herein refers to an aminocarbonyl as described above appended to the parent molecular moiety through an alkylene.

The term "aminocarbonylalkoxy" as used herein refers to  $\text{H}_2\text{N}-\text{C}(\text{O})-\text{O}-\text{R}_{55}-$ , wherein  $\text{R}_{55}$  is an alkyl radical group. Examples of aminocarbonylalkoxy include aminocarbonylmethoxy, aminocarbonylethoxy and the like.

30

The term "aryl" as used herein refers to a mono- or bicyclic carbocyclic ring system having one or two aromatic rings including, but not limited to, phenyl, naphthyl, tetrahydronaphthyl, indanyl, indenyl and the like. Aryl groups can be unsubstituted or substituted with one, two or three substituents independently  
5 selected from loweralkyl, halo, haloalkyl, haloalkoxy, hydroxy, oxo (=O), hydroxyalkyl, alkenyloxy, alkoxy, alkoxyalkoxy, alkoxycarbonyl, alkoxycarbonylalkenyl, (alkoxycarbonyl)thioalkoxy, thioalkoxy, alkylimino ( $R^*N=$  wherein  $R^*$  is a loweralkyl group), amino, alkylamino, alkylsulfonyl, dialkylamino, aminocarbonyl, aminocarbonylalkoxy, alkanoylamino, aryl, arylalkyl, arylalkoxy, 10 aryloxy, mercapto, cyano, nitro, mercapto, carboxy, carboxaldehyde, carboxamide, cycloalkyl, carboxyalkenyl, carboxyalkoxy, alkylsulfonylamino, cyanoalkoxy, (heterocyclic)alkoxy,  $-SO_3H$ , hydroxalkoxy, phenyl and tetrazolylalkoxy. Examples of substituted aryl include 3-chlorophenyl, 3-fluorophenyl, 4-chlorophenyl, 4-fluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluoro-phenyl, 4-methylsulfonylphenyl, 15 and the like.

The term "arylalkenyl" as used herein refers to an alkenylene to which is appended an aryl group, for example, phenylethenyl and the like.

The term "arylalkynyl" as used herein refers to an alkynyl radical to which is appended an aryl group, for example, phenylethynyl and the like

20 The term "arylalkoxy" as used herein refers to  $R_{42}O-$  wherein  $R_{42}$  is an arylalkyl group, for example, benzyloxy, and the like.

The term "arylalkyl" as used herein refers to an aryl group as previously defined, appended to a loweralkyl radical, for example, benzyl and the like.

25 The term "arylalkylamino" as used herein refers to an arylalkyl group as previously defined, appended to the parent molecular moiety through an amino group.

The term "arylamino" as used herein refers to  $R_{45}NH_2-$  wherein  $R_{45}$  is an aryl.

30 The term "arylcarbonylalkyl" as used herein refers to  $R_{45}C(O)R_{33}-$  wherein  $R_{45}$  is an aryl group and  $R_{33}$  is an alkylene group.

The term "aryloxy" as used herein refers to  $R_{45}O$ - wherein  $R_{45}$  is an aryl group, for example, phenoxy, and the like.

The term "carboxaldehyde" as used herein refers to a formaldehyde radical,  $-C(O)H$ .

5 The term "carboxamide" as used herein refers to  $-C(O)NH_2$ .

The term "carboxy" as used herein refers to a carboxylic acid radical,  $-C(O)OH$ .

The term "carboxyalkenyl" as used herein refers to a carboxy group as previously defined appended to an alkenyl radical as previously defined.

10 Examples of carboxyalkenyl include 2-carboxyethenyl, 3-carboxy-1-ethenyl and the like.

The term "carboxyalkyl" as used herein refers to a carboxy group as previously defined appended to an alkyl radical as previously defined. Examples of carboxyalkyl include 2-carboxyethyl, 3-carboxy-1-propyl and the like.

15 The term "carboxyalkoxy" as used herein refers to a carboxy group as previously defined appended to an alkoxy radical as previously defined. Examples of carboxyalkoxy include carboxymethoxy, carboxyethoxy and the like.

The term "cyano" as used herein refers a cyano ( $-CN$ ) group.

20 The term "cyanoalkyl" as used herein refers to a cyano ( $-CN$ ) group appended to the parent molecular moiety through an alkyl. Examples of cyanoalkyl include 3-cyanopropyl, 4-cyanobutyl, and the like.

The term "cyanoalkoxy" as used herein refers to a cyano ( $-CN$ ) group appended to the parent molecular moiety through an alkoxy radical. Examples of cyanoalkoxy include 3-cyanopropoxy, 4-cyanobutoxy and the like.

25 The term "cycloalkyl" as used herein refers to an aliphatic ring system having 3 to 10 carbon atoms and 1 to 3 rings including, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl, and the like. Cycloalkyl groups can be unsubstituted or substituted with one, two or three substituents independently selected from hydroxy, halo, oxo ( $=O$ ), alkylimino ( $R^*N=$  wherein  $R^*$  is a loweralkyl group), amino, 30 alkylamino, dialkylamino, alkoxy, alkoxyalkoxy, alkoxy carbonyl, thioalkoxy,



haloalkyl, mercapto, carboxy, carboxaldehyde, carboxamide, cycloalkyl, aryl, arylalkyl,  $-\text{SO}_3\text{H}$ , nitro, cyano and loweralkyl.

5 The term "cycloalkenyl" as used herein refers to an aliphatic ring system having 3 to 10 carbon atoms and 1 to 3 rings containing at least one double bond in the ring structure. Cycloalkenyl groups can be unsubstituted or substituted with one, two or three substituents independently selected hydroxy, halo, oxo ( $=\text{O}$ ), alkylimino ( $\text{R}^*\text{N}=$  wherein  $\text{R}^*$  is a loweralkyl group), amino, alkylamino, dialkylamino, alkoxy, alkoxyalkoxy, alkoxycarbonyl, thioalkoxy, haloalkyl, mercapto, carboxy, carboxaldehyde, carboxamide, cycloalkyl, aryl, arylalkyl,  $-\text{SO}_3\text{H}$ , nitro, 10 cyano and loweralkyl.

The term "cycloalkylalkyl" as used herein refers to a cycloalkyl group appended to a loweralkyl radical, including but not limited to cyclohexylmethyl.

The term "cycloalkenylalkyl" as used herein refers to a cycloalkenyl group appended to a loweralkyl radical, including but not limited to cyclohexenylmethyl.

15 The term "dialkylamino" as used herein refers to  $(\text{R}_{56})(\text{R}_{57})\text{N}-$  wherein  $\text{R}_{56}$  and  $\text{R}_{57}$  are independently selected from loweralkyl, for example diethylamino, methyl propylamino, and the like.

The term "diarylamino" as used herein refers to  $(\text{R}_{45})(\text{R}_{46})\text{N}-$  wherein  $\text{R}_{45}$  and  $\text{R}_{46}$  are independently aryl, for example diphenylamino and the like.

20 The term "halo" or "halogen" as used herein refers to I, Br, Cl or F.

The term "haloalkyl" as used herein refers to an alkyl radical, as defined above, which has at least one halogen substituent, for example, chloromethyl, fluoroethyl, trifluoromethyl or pentafluoroethyl, 2,3-difluoropentyl, and the like.


25 The term "haloalkenyl" as used herein refers to an alkenyl radical has at least one halogen substituent.

The term "haloalkynyl" as used herein refers to an alkynyl radical has at least one halogen substituent.

30 The term "haloalkoxy" as used herein refers to an alkoxy radical as defined above, bearing at least one halogen substituent, for example, 2-fluoroethoxy, 2,2,2-trifluoroethoxy, trifluoromethoxy,

2,2,3,3,3-pentafluoropropoxy and the like.

The term "heterocyclic ring" or "heterocyclic" or "heterocycle" as used herein refers to any 3- or 4-membered ring containing a heteroatom selected from oxygen, nitrogen and sulfur; or a 5-, 6- or 7-membered ring containing one, two or three nitrogen atoms; one oxygen atom; one sulfur atom; one nitrogen and one sulfur atom; one nitrogen and one oxygen atom; two oxygen atoms in non-adjacent positions; one oxygen and one sulfur atom in non-adjacent positions; or two sulfur atoms in non-adjacent positions. Examples of heterocycles include, but are not limited to, thiophene, pyrrole, and furan. The 5-membered ring has 0-2 double bonds and the 6- and 7-membered rings have 0-3 double bonds. The nitrogen heteroatoms can be optionally quaternized. The term "heterocyclic" also includes bicyclic groups in which any of the above heterocyclic rings is fused to a benzene ring or a cycloalkane ring or another heterocyclic ring (for example, indolyl, dihydroindolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, decahydroquinolyl, decahydroisoquinolyl, benzofuryl, dihydrobenzofuryl or benzothienyl and the like). Heterocyclics include: aziridinyl, azetidiny, pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, imidazolyl, imidazoliny, imidazolidinyl, pyridyl, piperidinyl, homopiperidinyl, pyrazinyl, piperazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiomorpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, oxetanyl, furyl, tetrahydrofuranly, thienyl, thiazolidinyl, isothiazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, thiadiazolyl, pyrrolyl, pyrimidyl and benzothienyl.



Heterocyclics also include compounds of the formula  $\text{X}^* \text{---} \text{O} \text{---} \text{O} \text{---} \text{C}(\text{R}'')_2$  where  $\text{X}^*$  is  $-\text{CH}_2-$  or  $-\text{O}-$  and  $\text{Y}^*$  is  $-\text{C}(\text{O})-$  or  $-\text{C}(\text{R}'')_2-$ , where  $\text{R}''$  is hydrogen or  $\text{C}_1$ - $\text{C}_4$ -alkyl and  $v$  is 1, 2 or 3 such as 1,3-benzodioxolyl, 1,4-benzodioxanyl and the like. Heterocyclics also include bicyclic rings such as quinuclidinyl and the like.

Heterocyclics can be unsubstituted or be substituted with one, two, or three substituents independently selected from hydroxy, halo, oxo ( $=\text{O}$ ), alkylimino ( $\text{R}^*\text{N}=\text{O}$ ) wherein  $\text{R}^*$  is a loweralkyl group), amino, alkylamino, dialkylamino, alkoxy,

alkoxyalkoxy, alkoxycarbonyl, thioalkoxy, haloalkyl, mercapto, carboxy, carboxaldehyde, carboxamide, cycloalkyl, aryl, arylalkyl, -SO<sub>3</sub>H, nitro, cyano and loweralkyl. In addition, nitrogen containing heterocycles can be N-protected.

5 The term "heterocyclic(alkoxy)" as used herein refers to a heterocyclic group as defined above appended to an alkoxy radical as defined above. Examples of (heterocyclic)alkoxy include 4-pyridylmethoxy, 2-pyridylmethoxy and the like.

The term "heterocyclic(alkyl)" as used herein refers to a heterocyclic group as defined above appended to the parent molecular moiety through a loweralkyl radical as defined above.

10 The term "heterocyclic(oxy)" as used herein refers to a heterocyclic group as defined above appended to the parent molecular moiety through an oxygen. Examples of (heterocyclic)oxy include 4-pyridyloxy, 2-pyridyloxy and the like.

The term "hydroxy" as used herein refers to -OH.

15 The term "hydroxyalkoxy" as used herein refers to an alkoxy radical as previously defined to which is appended a hydroxy (-OH) group. Examples of hydroxyalkoxy include 3-hydroxypropoxy, 4-hydroxybutoxy and the like.

The term "hydroxyalkyl" as used herein refers to a loweralkyl radical to which is appended a hydroxy group.

The term "mercapto" or "thiol" as used herein refers to -SH.

20 The term "nitro" as used herein refers to -NO<sub>2</sub>.

The term "thioalkoxy" as used herein refers to R<sub>70</sub>S- wherein R<sub>70</sub> is alkoxy. Examples of thioalkoxy include, but are not limited to, methylthio, ethylthio and the like.

25 The compounds of the present invention can be used in the form of salts derived from inorganic or organic acids. These salts include but are not limited to the following: acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, cyclopentanepropionate, dodecylsulfate, ethanesulfonate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, 30 hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactate, maleate,

methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as loweralkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides, and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl, and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides, and others. Water or oil-soluble or dispersible products are thereby obtained.

10        Examples of acids which may be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

15        Basic addition salts can be prepared *in situ* during the final isolation and purification of the compounds of formula (I), or separately by reacting a carboxylic acid function with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia, or an organic primary, secondary or tertiary amine. Such pharmaceutically acceptable salts include, but are not limited to, cations based on the alkali and alkaline earth metals, 20        such as sodium, lithium, potassium, calcium, magnesium, aluminum salts and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. Other representative organic amines useful for the 25        formation of base addition salts include diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like.

30        The term "pharmaceutically acceptable ester" as used herein refers to esters which hydrolyze *in vivo* and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanolic, alkenolic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6

carbon atoms. Examples of particular esters includes formates, acetates, propionates, butyates, acrylates and ethylsuccinates.

The term "pharmaceutically acceptable prodrug" as used herein refers to those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "prodrug" refers to compounds that are rapidly transformed *in vivo* to provide the parent compound having the above formula, for example by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, Vol. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press, 1987, both of which are incorporated herein by reference.

As used throughout this specification and the appended claims, the term metabolically cleavable group denotes a moiety which is readily cleaved *in vivo* from the compound bearing it, wherein said compound, after cleavage remains or becomes pharmacologically active. Metabolically cleavable groups form a class of groups reactive with the carboxyl group of the compounds of this invention are well known to practitioners of the art. They include, but are not limited to groups such as, for example, alkanoyl, such as acetyl, propionyl, butyryl, and the like; unsubstituted and substituted aroyl, such as benzoyl and substituted benzoyl; alkoxycarbonyl, such as ethoxycarbonyl; trialkylsilyl, such as trimethyl- and triethysilyl; monoesters formed with dicarboxylic acids, such as succinyl, and the like. Because of the ease with which the metabolically cleavable groups of the compounds of this invention are cleaved *in vivo*, the compounds bearing such groups act as pro-drugs of other prostaglandin biosynthesis inhibitors. The compounds bearing the metabolically cleavable groups have the advantage that they may exhibit improved bioavailability as a result of enhanced solubility and/or rate of absorption conferred upon the parent compound by virtue of the presence of the metabolically cleavable group.

Asymmetric centers may exist in the compounds of the present invention. The present invention contemplates the various stereoisomers and mixtures thereof. Individual stereoisomers of compounds of the present invention are made by synthesis from starting materials containing the chiral centers or by preparation  
5 of mixtures of enantiomeric products followed by separation as, for example, by conversion to a mixture of diastereomers followed by separation by recrystallization or chromatographic techniques, or by direct separation of the optical enantiomers on chiral chromatographic columns. Starting compounds of particular stereochemistry are either commercially available or are made by the methods  
10 detailed below and resolved by techniques well known in the organic chemical arts.

The following examples illustrate the process of the invention, without limitation.

#### EXAMPLE 1

##### 15 2-(4-Fluorophenyl)-4-methylthio-*trans*-cinnamic acid

To a stirred solution of 4-fluorophenylacetic acid (48.8 g, 0.31 mol.) and 4-(methylthio)benzaldehyde (50 g, 0.31 mol.) in acetic anhydride (50 mL) was added sodium methoxide (18.8 g, 0.33 mol.). The solution was stirred at reflux for 18 hours. The reaction mixture was allowed to cool to room temperature and a  
20 yellow precipitate formed. The yellow precipitate was filtered, stirred in water (550 mL) for 3 hours, and refiltered. The product was purified by recrystallization (ethanol) to provide yellow crystals (yield: 51.55 g; 57%). M.p. = 163-167°C.

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ 2.42 (s, 3 H), 6.97 (d, J = 9 Hz, 2 H), 7.08 (d, J = 9 Hz, 2 H), 7.2 (d, J = 7 Hz, 4 H), 7.72 (s, 1 H), 12.68 (bs, 1 H).

25 MS (DCI/NH<sub>3</sub>) m/e 289 (M + H)<sup>+</sup>.

#### EXAMPLE 2

##### 4-Fluorophenyl-4'-methylthiobenzylketone

A solution of 2-(4-fluorophenyl)-4-methylthio-*trans*-cinnamic acid, prepared in  
30 Example 1, (51.48 g, 0.179 mol.) in thionyl chloride (115 mL) was heated at reflux

for 1.5 hours and then stirred at room temperature for an additional 18 hours. The reaction mixture was concentrated *in vacuo*, dissolved in acetone, and added dropwise to a solution of sodium azide (12.8 g, 0.197 mol.) in water (95 mL) maintained at 0°C. The mixture was allowed to warm to room temperature, diluted with water (730 mL), and extracted with toluene (1 L). The organic extract was washed with brine, dried over MgSO<sub>4</sub>, and filtered. The filtrate was heated at reflux for 1.5 hours, concentrated *in vacuo*, and dissolved in acetic acid:water (127 mL:63 mL). The solution was heated at reflux for 2 hours, allowed to cool to room temperature, diluted with water (180 mL), whereafter a yellow solid precipitated, and was filtered. The yellow precipitate was purified by recrystallization (ethanol) to provide yellow crystals (yield: 27.85 g; 59%).

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.05 (s, 3 H), 5.43 (s, 2 H), 6.95 (m, 3 H), 7.1 (m, 2 H), 7.3 (d, J = 9 Hz, 2 H), 7.37 (m, 3 H), 7.55 (m, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 435 (M + H)<sup>+</sup>, 452 (M + NH<sub>4</sub>)<sup>+</sup>.

Elemental analysis, calculated for C<sub>24</sub>H<sub>19</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 66.34; H, 4.40; N, 6.44.  
Found: C, 66.21; H, 4.35; N, 6.43.

### EXAMPLE 3

#### 3-(4-Fluorobenzoyl)-3-(4-methylthiophenyl)propionic acid, ethyl ester

A solution of 4-fluorophenyl-4'-methylthiobenzylketone, prepared in Example 2, (10 g, 38.45 mmol.) in THF (300 mL), was prepared and maintained at -78°C. A 1M THF solution of sodium bis(trimethylsilyl)amide (38.45 mL, 38.45 mmol.) was added dropwise. The reaction mixture was stirred at -78°C for 45 minutes and ethyl bromoacetate (4.26 mL, 38.45 mmol.) was added dropwise maintaining the temperature at -78°C. The mixture was allowed to warm to room temperature and stirred for 18 hours. The solution was acidified with a 10 % citric acid solution, extracted with ether (3 x 50 mL), washed with brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. The residue was purified by chromatography (silica gel, 5 % ethyl acetate/hexane) to provide an oil (yield: 8.7 g; 65%).

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ 1.1 (s, 3 H), 2.41 (s, 3 H), 2.66 & 2.71 (2d, J = 5 Hz, 1 H), 3.14 & 3.20 (2d, J = 5 Hz, 1 H), 4.02 (q, J = 7 Hz, 2 H), 5.19 & 5.23 (2d, J = 5 Hz, 1 H), 7.18 (d, J = 9 Hz, 2 H), 7.29 (m, 4 H), 8.1 (m, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 347 (M + H)<sup>+</sup>.

#### EXAMPLE 4

3-(4-Fluorobenzoyl)-3-(4-methylsulfonylphenyl)propionic acid, ethyl ester

To a stirred solution of ethyl 3-(4-fluorobenzoyl)-3-(4'-methylthiophenyl)propionate,  
5 prepared in Example 3 (8.7 g, 25.11 mmol.), in CH<sub>2</sub>Cl<sub>2</sub> (450 mL) at 0°C was added  
3-chloroperoxybenzoic acid (17.3 g, 50.22 mmol.). The reaction mixture was  
warmed to room temperature and stirred for 45 minutes. The solution was  
quenched with a saturated aqueous solution of sodium sulfite. The organic layer  
was washed with 1N aqueous NaOH (2 x 50 mL), brine, and dried over MgSO<sub>4</sub>.  
10 The solution was concentrated *in vacuo* to provide an oil (yield: 9.45 g; 99%).

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ 1.1 (t, J = 7 Hz, 3 H), 2.76 & 2.82 (2d, J = 5  
Hz, 1 H), 3.17 (s, 3 H), 3.18-3.32 (m, 1 H), 4.03 (q, J = 7 Hz, 2 H), 5.42 & 5.45 (2d, J  
= 5 Hz, 1 H), 7.32 (t, J = 9 Hz, 3 H), 7.65 (d, J = 9 Hz, 2 H), 7.84 (d, J = 9 Hz, 2 H),  
8.16 (dd, J = 9 Hz, 7 Hz, 2 H).

15 MS (DCI/NH<sub>3</sub>) m/e 379 (M + H)<sup>+</sup>.

#### EXAMPLE 5

5-(4-Methylsulfonylphenyl)-6-(4-fluorophenyl)-4,5-dihydro-3(2H)-pyridazinone

To a stirred solution of ethyl 3-(4-fluorobenzoyl)-3-(4-methylsulfonylphenyl)-  
20 propionate, prepared in Example 3 (9.7 g, 26.99 mmol.), in ethanol (140 mL) was  
added hydrazine monohydrate (40 mL, 0.824 mol.). The reaction vessel was  
equipped with a soxhlet extractor, heated at reflux for 7 hours. The reaction  
mixture was concentrated *in vacuo*, quenched with ice water (100 mL) and  
concentrated HCl (10 mL), extracted with ethyl acetate (3 x 50 mL). The combined  
25 extracts were washed with brine, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*.  
The product was purified by recrystallization (MeOH) to provide a yellow solid  
(yield: 3.79 g; 41%). M.p. = 240-241°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 2.8 (m, 1 H), 3.05 (s, 3 H), 3.1 (m, 1 H), 4.58 (m,  
1 H), 7.05 (t, J = 9 Hz, 2 H), 7.42 (d, J = 9 Hz, 2 H), 7.65 (m, 2 H), 7.91 (d, J = 9 Hz, 2  
30 H), 8.88 (bs, 1 H).

MS (DCI/NH<sub>3</sub>) m/e 347 (M + H)<sup>+</sup>.



**EXAMPLE 6****5-(4-Methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

To a stirred solution of 5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-4,5-dihydro-3(2H)-pyridazinone, prepared in Example 5 (2.6 g, 7.51 mmol.), in glacial acetic acid (250 mL) maintained at 95°C was added bromine (0.386 mL, 7.51 mmol.). The reaction mixture was stirred at 95°C for 30 minutes, concentrated *in vacuo*, quenched with water, and extracted with ethyl acetate (2 x 25 mL). The combined extracts were washed with brine, dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and filtered to provide a yellow solid (yield: 2.13 g; 82 %). M.p. = 260-262°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.08 (s, 3 H), 6.98 (m, 3 H), 7.13 (m, 2 H), 7.35 (d, J = 9 Hz, 2 H), 7.92 (d, J = 9 Hz, 2 H), 11.1 (bs, 1 H).

MS (DCI/NH<sub>3</sub>) m/e 345 (M + H)<sup>+</sup>.

**EXAMPLE 7****2-Benzyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

A solution of 5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone, prepared in Example 6 (600 mg, 1.74 mmol.), K<sub>2</sub>CO<sub>3</sub> (264 mg, 1.91 mmol.), benzyl bromide (0.207 mL, 1.74 mmol.) and NaI (about 0.1 eq., catalytic) in about 40 mL of dimethylformamide was stirred at room temperature for 18 hours. The reaction mixture was quenched with 2N HCl, and extracted with ethyl acetate (2 x 20 mL). The combined extracts were washed with brine, water, and dried over MgSO<sub>4</sub>. The dried extracts were concentrated *in vacuo* and the residue purified by recrystallization (MeOH) to provide a white solid (yield: 601 mg; 79%). M.p. = 172-174°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.05 (s, 3 H), 5.43 (s, 2 H), 6.95 (m, 3 H), 7.1 (m, 2 H), 7.3 (d, J = 9 Hz, 2 H), 7.37 (m, 3 H), 7.55 (m, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 435 (M + H)<sup>+</sup>, 452 (M + NH<sub>4</sub>)<sup>+</sup>.

Elemental analysis, , calculated for C<sub>24</sub>H<sub>19</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 66.34; H, 4.40; N, 6.44.

Found: C, 66.21; H, 4.35; N, 6.43.

**EXAMPLE 8****2-Methyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting methyl iodide for benzyl bromide. M.p. = 181-182°C.

5  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  3.08 (s, 3 H), 3.92 (s, 3 H), 6.98 (m, 3 H), 7.13 (m, 2 H), 7.32 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/ $\text{NH}_3$ ) m/e 359 (M + H) $^+$ , 376 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{18}\text{H}_{15}\text{FN}_2\text{O}_3\text{S}$ : C, 60.32; H, 4.21; N, 7.81.

Found: C, 60.26; H, 3.93; N, 7.81.

10

**EXAMPLE 9****2-Ethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting ethyl bromide for benzyl bromide. M.p. = 177-179°C.

15  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  1.47 (t, J = 7 Hz, 3 H), 3.05 (s, 3 H), 4.33 (q, J = 7 Hz, 2 H), 6.96 (m, 3 H), 7.13 (m, 2 H), 7.32 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/ $\text{NH}_3$ ) m/e 373 (M + H) $^+$ , 390 (M +  $\text{NH}_4$ ) $^+$ .

Elemental analysis, calculated for  $\text{C}_{19}\text{H}_{17}\text{FN}_2\text{O}_3\text{S} \cdot 0.25\text{M H}_2\text{O}$ :

C, 60.54; H, 4.67; N, 7.43;

20

Found: C, 60.40; H, 4.55; N, 7.43.

**EXAMPLE 10****2-(4-Fluorobenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

25 The title compound was prepared according to the method of Example 7, substituting 4-fluorobenzyl bromide for benzyl bromide. M.p. = 150-151°C.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  3.05 (s, 3 H), 5.4 (s, 2 H), 6.92-7.14 (m, 7 H), 7.3 (d, J = 9 Hz, 2 H), 7.52 (m, 2 H), 7.9 (d, J = 9 Hz, 2 H).

30 MS (DCI/ $\text{NH}_3$ ) m/e 453 (M + H) $^+$ , 470 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{24}\text{H}_{18}\text{F}_2\text{N}_2\text{O}_3\text{S}$ : C, 63.70; H, 4.00; N, 6.19.

Found: C, 63.46; H, 3.86; N, 5.92.

### EXAMPLE 11

**2-(n-Butyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting n-butyl bromide for benzyl bromide. M.p. = 111-113°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.0 (t, J = 7 Hz, 3 H), 1.47 (m, 2 H), 1.9 (m, 2 H), 3.08 (s, 3 H), 4.28 (t, J = 7 Hz, 2 H), 6.96 (m, 3 H), 7.14 (m, 2 H), 7.33 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 401 (M + H)<sup>+</sup>, 418 (M + NH<sub>4</sub>)<sup>+</sup>.

Elemental Analysis: calculated for  $C_{21}H_{21}FN_2O_3S$ : C, 62.98; H, 5.28; N, 6.99.

Found: C, 62.95; H, 4.67; N, 6.90.

### EXAMPLE 12

2-(4-Fluoro- $\alpha$ -methylbenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting 4-fluoro- $\alpha$ -methyl benzyl bromide (synthesized from 4-fluoro- $\alpha$ -methyl benzyl alcohol) for benzyl bromide. M.p. = 185-186°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 1.84 (d, J = 7 Hz, 3 H), 3.07 (s, 3 H), 6.43 (q, J = 7 Hz, 1 H), 6.91-7.11 (m, 7 H), 7.31 (d, J = 9 Hz, 2 H), 7.52 (m, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 467 (M + H)<sup>+</sup>, 484 (M + NH<sub>4</sub>)<sup>+</sup>.

**Elemental Analysis:** calculated for  $C_{25}H_{20}F_2N_2O_3S$ :

C, 64.36; H, 4.32; N, 6.00.

Found: C, 64.38; H, 4.07; N, 5.93.

### EXAMPLE 13

**2-(n-Propyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting propyl bromide for benzyl bromide. M.p. = 155-157°C.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  1.04 (t, J = 7 Hz, 3 H), 1.94 (m, 2 H), 3.08 (s, 3 H), 4.23 (t, J = 7 Hz, 2 H), 6.96 (m, 3 H), 7.13 (m, 2 H), 7.33 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H);

MS (DCI/ $\text{NH}_3$ ) m/e 387 (M + H) $^+$ , 404 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{20}\text{H}_{19}\text{FN}_2\text{O}_3\text{S}$ : C, 62.16; H, 4.95; N, 7.24.

Found: C, 62.15; H, 4.78; N, 7.22.

#### EXAMPLE 14

##### 2-(n-Pentyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting n-pentyl bromide for benzyl bromide. M.p. = 65-67°C.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  0.93 (m, 3 H), 1.2 (m, 4 H), 1.9 (m, 2 H), 3.08 (s, 3 H), 4.28 (t, J = 7 Hz, 2 H), 6.97 (m, 3 H), 7.14 (m, 2 H), 7.33 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/ $\text{NH}_3$ ) m/e 415 (M + H) $^+$ , 432 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{22}\text{H}_{23}\text{FN}_2\text{O}_3\text{S}\cdot 0.75\text{M H}_2\text{O}$ :

C, 61.73; H, 5.76; N, 6.54.

Found: C, 61.97; H, 6.00; N, 6.36.

#### EXAMPLE 15

##### 2-Cyclohexylmethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting cyclohexylmethyl bromide for benzyl bromide. M.p. = 175-176°C.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  1.03-1.38 (m, 5 H), 1.63-1.8 (m, 5 H), 2.05 (m, 1 H), 3.08 (s, 3 H), 4.13 (d, J = 7 Hz, 2 H), 6.97 (m, 3 H), 7.14 (m, 2 H), 7.33 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS (DCI/ $\text{NH}_3$ ) m/e 441 (M + H) $^+$ , 458 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{24}\text{H}_{25}\text{FN}_2\text{O}_3\text{S}\cdot 0.25\text{M H}_2\text{O}$ :

C, 64.77; H, 5.77; N, 6.29.

Found: C, 64.63; H, 5.85; N, 6.16.

### EXAMPLE 16

#### 2-Phenacyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

5 The title compound was prepared according to the method of Example 7, substituting 2-bromoacetophenone for benzyl bromide. M.p. = 178-180°C.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  3.08 (s, 3 H), 5.75 (s, 2 H), 6.96 (t, J = 9 Hz, 2 H), 7.03 (s, 1 H), 7.15 (dd, J = 9 Hz, 7 Hz, 2 H), 7.37 (d, J = 9 Hz, 2 H), 7.55 (t, J = 7 Hz, 2 H), 7.68 (t, J = 7 Hz, 1 H), 7.94 (d, J = 9 Hz, 2 H), 8.05 (d, J = 7 Hz, 2 H).

10 MS (DCI/ $\text{NH}_3$ ) m/e 463 (M + H) $^+$ , 480 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{25}\text{H}_{19}\text{FN}_2\text{O}_4\text{S} \cdot 0.75 \text{ M H}_2\text{O}$ :

C, 63.08; H, 4.34; N, 5.88.

Found: C, 63.01; H, 4.12; N, 5.85.

### 15 EXAMPLE 17

#### 2-Propargyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting propargyl bromide for benzyl bromide. M.p. = 168-169°C.

20  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  2.4 (t, J = 3 Hz, 1 H), 3.08 (s, 3 H), 5.05 (d, J = 3 Hz, 2 H), 6.97 (t, J = 9 Hz, 2 H), 6.99 (s, 1 H), 7.16 (dd, J = 9 Hz, 7 Hz, 2 H), 7.34 (d, J = 9 Hz, 2 H), 7.91 (d, J = 9 Hz, 2 H).

MS (DCI/ $\text{NH}_3$ ) m/e 383 (M + H) $^+$ , 400 (M +  $\text{NH}_4$ ) $^+$ .

Elemental Analysis: calculated for  $\text{C}_{20}\text{H}_{15}\text{FN}_2\text{O}_3\text{S} \cdot 0.25 \text{ M H}_2\text{O}$ :

C, 62.08; H, 4.03; N, 7.24.

25 Found: C, 62.15; H, 4.18; N, 7.05.

### EXAMPLE 18

#### 2-Cyclohexyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

30 The title compound was prepared according to the method of Example 7, substituting cyclohexyl bromide for benzyl bromide. M.p. = 86-90 °C.

$^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$  1.5 (m, 10 H), 3.24 (s, 3 H), 4.85 (m, 1 H), 7.04 (s, 1 H), 7.19 (m, 4 H), 7.46 (d,  $J = 9$  Hz, 2 H), 7.87 (d,  $J = 9$  Hz, 2 H).

MS (DCI/ $\text{NH}_3$ )  $m/e$  427 ( $M + H$ ) $^+$ .

Elemental Analysis calculated. for  $\text{C}_{23}\text{H}_{23}\text{FN}_2\text{O}_3\text{S}$ : C, 64.77; H, 5.43; N, 6.56.

Found: C, 64.52; H, 5.4; N, 6.38.

#### EXAMPLE 19

##### 2-(2-Butyn-1-yl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7,

substituting 1-bromo-2-butyne for benzyl bromide. M.p. = 81-82 °C.

$^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$  1.81 (t,  $J = 3$  Hz, 3 H), 3.24 (s, 3 H), 4.93 (d,  $J = 3$ , 2 H), 7.10 (s, 1 H), 7.19 (m, 4 H), 7.48 (d,  $J = 9$  Hz, 2 H), 7.87 (d,  $J = 9$  Hz, 2 H).

MS (DCI/ $\text{NH}_3$ )  $m/e$  397 ( $M+H$ ) $^+$ .

Elemental analysis, calculated for  $\text{C}_{21}\text{H}_{17}\text{FN}_2\text{O}_3\text{S}$ :

C, 63.62; H, 4.32; N, 7.06.

Found: C, 64.22; H, 4.36; N, 6.61.

#### EXAMPLE 20

##### 2-(Cyclobutylmethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7,

substituting chloromethylcyclobutane for benzyl bromide. M.p. = 74-76 °C.

$^1\text{H}$  NMR (300 MHz,  $d_6$ -DMSO)  $\delta$  1.98 (m, 4 H), 2.05 (m, 2 H), 2.84 (p,  $J = 7.5$  Hz, 1 H), 3.24 (s, 3 H), 4.22 (d,  $J = 7.5$  Hz, 2 H), 7.05 (s, 1 H), 7.16 (m, 4 H), 7.45 (d,  $J = 9$  Hz, 2 H), 7.87 (d,  $J = 9$  Hz, 2 H).

MS (DCI/ $\text{NH}_3$ )  $m/e$  413 ( $M+H$ ) $^+$ .

Elemental analysis, calculated for  $\text{C}_{22}\text{H}_{21}\text{FN}_2\text{O}_3\text{S}$ : C, 64.06; H, 5.13; N, 6.79.

Found: C, 64.37; H, 5.26; N, 6.73.

**EXAMPLE 21****2-(3-Methylbuten-2-yl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting 4-bromo-2-methyl-2-butene for benzyl bromide. M.p. = 71-72 °C.

<sup>1</sup>H NMR (300 MHz, d<sub>6</sub>-DMSO) δ 1.72 (s, 3 H), 1.79 (s, 3 H), 3.24 (s, 3 H), 4.76 (d, J = 7.5 Hz, 2 H), 5.42 (m, 1 H), 7.04 (s, 1 H), 7.17 (m, 4 H), 7.46 (d, J = 9 Hz, 2 H), 7.87 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 413 (M+H)<sup>+</sup>.

Elemental analysis, calculated for C<sub>22</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 64.06; H, 5.13; N, 6.79.

Found: C, 64.34; H, 4.87; N, 6.51.

**EXAMPLE 22****2-(2,2,2-Trifluoroethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound is prepared according to the method of Example 7, substituting 2-iodo-1,1,1-trifluoroethane for benzyl bromide. M.p. = 177-179°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.08 (s, 3 H), 4.90 (q, 2 H), 6.98 (t, J = 9 Hz, 2 H), 7.01 (s, 1 H), 7.13 (dd, J = 9 Hz, 7 Hz, 2 H), 7.34 (d, J = 9 Hz, 2 H), 7.92 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 427 (M + H)<sup>+</sup>, 444 (M + NH<sub>4</sub>)<sup>+</sup>.

Elemental analysis, calculated for C<sub>19</sub>H<sub>14</sub>F<sub>3</sub>N<sub>2</sub>O<sub>3</sub>S (0.25 hydrate):

C, 52.96; H, 3.39; N, 6.50.

Found: C, 52.89; H, 3.35; N, 6.27.

**EXAMPLE 23****2-(Cyclopropylmethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone**

The title compound was prepared according to the method of Example 7, substituting cyclopropylmethyl bromide for benzyl bromide. M.p. = 130-132°C.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) 0.5 (m, 2 H), 0.6 (m, 2 H), 1.46 (m, 1 H), 3.08 (s, 3 H), 4.15 (d, J = 7 Hz, 2 H), 6.97 (s, 1 H), 6.98 (d, J = 9 Hz, 2 H), 7.14 (dd, J = 9 Hz, 7 Hz, 2 H), 7.35 (d, J = 9 Hz, 2 H), 7.9 (d, J = 9 Hz, 2 H).

MS m/e 399 (M + H)<sup>+</sup>, 416 (M + NH<sub>4</sub>)<sup>+</sup>.

Elemental analysis, calculated for C<sub>21</sub>H<sub>19</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 63.30; H, 4.80; N, 7.03.

Found: C, 63.13; H, 4.75; N, 6.94.

#### EXAMPLE 24

##### 2-Cyclopentylmethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting chloromethylcyclopentane for benzyl bromide. M.p. = 76-77°C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 300 MHz) δ 1.38 (m, 2 H), 1.6 (m, 7 H), 3.23 (s, 3 H), 4.11 (d, J = 7.5 Hz, 2 H), 7.05 (s, 1 H), 7.17 (m, 4 H), 7.47 (d, J = 9 Hz, 2 H), 7.87 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 427 (M+H)<sup>+</sup>.

Elemental analysis, calculated for C<sub>23</sub>H<sub>23</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 64.77; H, 5.43; N, 6.56.

Found: C, 63.96; H, 5.11; N, 6.44.

#### EXAMPLE 25

##### 2-Cyclopentyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone

The title compound was prepared according to the method of Example 7, substituting chlorocyclopentane for benzyl bromide. M.p. = 191-192°C.

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO, 300 MHz) δ 1.65 (m, 2 H), 1.85 (m, 4 H), 2.05 (m, 2 H), 3.24 (s, 3 H), 5.39 (p, J = 7.5 Hz, 1 H), 7.02 (s, 1 H), 7.18 (m, 4 H), 7.46 (d, J = 9 Hz, 2 H), 7.89 (d, J = 9 Hz, 2 H).

MS (DCI/NH<sub>3</sub>) m/e 413 (M+H)<sup>+</sup>.

Elemental analysis, calculated for C<sub>22</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>3</sub>S: C, 64.06; H, 5.13; N, 6.79.



Found: C, 64.14; H, 5.11; N, 6.69

## Prostaglandin Inhibition Determination

### Compound Preparation and Administration

For oral administration, test compounds were suspended on the day of use  
5 in 100% polyethyleneglycol (PEG 400) with a motorized homogenizer equipped  
with a Teflon-coated pestle (TRI-R Instrument, Jamaica, NY).

To compare the mean responses of the treatment groups, analysis of  
variance was applied. Percent inhibition values were determined by comparing  
the individual treatment mean values to the mean of the control group. Linear  
10 regression was used to estimate  $IC_{50}$ 's/ $ED_{50}$ 's in appropriate assays.

### EIA Determination of Prostaglandins

EIA reagents for prostaglandin determination were purchased from  
Perseptive Diagnostics, (Cambridge, MA).  $PGE_2$  levels in lavage fluids were  
determined after the samples were dried under nitrogen and reconstituted with  
15 assay buffer.  $PGE_2$  levels in enzyme assays or cell culture media were measured  
against standards prepared in the same milieu. The immunoassays were  
conducted as recommended by the manufacturer. The EIA was conducted in 96  
well microtiter plates (Nunc Roskilde, Denmark) and optical density was measured  
using a microplate reader (Vmax, Molecular Devices Corp., Menlo Park, CA).

### 20 Recombinant Human PGHS-1 and PGHS-2 Enzyme Assays

Inhibition of prostaglandin biosynthesis *in vitro* was evaluated using  
recombinant human Cox-1 (r-hu Cox1) and Cox-2 (r-hu Cox2) enzyme assays.  
Representative compounds dissolved in DMSO (3.3% v/v) were preincubated with  
microsomes from recombinant human Cox-1 or Cox-2 expressed in the  
25 baculovirus/Sf9 cell system (Gierse, J. K., Hauser, S. D., Creely, D. P., Koboldt, C.,  
Rangwala, S., H., Isakson, P. C., and Seibert, K. Expression and selective  
inhibition of the constitutive and inducible forms of cyclooxygenase, *Biochem J.*  
1995, 305: 479.), together with the cofactors phenol (2 mM) and hematin (1  $\mu$ M) for  
60 minutes prior to the addition of 10  $\mu$ M arachidonic acid. The reaction was

- allowed to run for 2.5 minutes at room temperature prior to quenching with HCl and neutralization with NaOH. PGE<sub>2</sub> production in the presence and absence of the drug was determined by EIA analysis. The EIA was conducted in 96 well microtiter plates (Nunc Roskilde, Denmark) and optical density was measured using a microplate reader (Vmax, Molecular Devices Corp., Menlo Park, CA). EIA reagents for prostaglandin determination were purchased from Perseptive Diagnostics (Cambridge, MA). PGE<sub>2</sub> levels were measured against standards prepared in the same milieu. The immunoassays were conducted as recommended by the manufacturer.
- 10 The data illustrating the inhibition of prostaglandin biosynthesis *in vitro* by compounds of this invention is shown in Table 1. The compounds are designated by the Example Number. Column 2 shows Cox-1 percent inhibition at the particular micromolar dose level. Column 3 shows Cox-2 percent inhibition at the particular nanomolar dose level. Values for Cox-2 inhibition that are parenthetical indicate IC<sub>50</sub> values.

TABLE 1

Example No.	RHUCX1 ( $\mu$ M)	RHUCX2 (n M)
7	4@100	(20)
10	2@100	65@100
11	19@100	51@100
12	8@100	(740)
14	23@100	66@100
15	52@100	(5)
17	0 @ 100	(850)
18	16@100	49 @ 100
19	3 @100	66 @1000
21	62 @ 100	92 @ 100
22	0@100	46@1000

23	24@100	(250)
24	23 @ 100	87 @100
25	7 @ 100	42 @ 100

### IL-1 $\beta$ Induced PGE<sub>2</sub> Production in WISH Cells

Human amnionic WISH cells were grown to 80% confluence in 48 well plates. Following removal of the growth medium and two washings with Gey's  
 5 Balanced Salt Solution, 5 ng IL-1 $\beta$ /ml (UBI, Lake Placid, NY) was added to the cells with or without test compound in DMSO (0.01% v/v) in Neuman-Tytell Serumless Medium (GIBCO, Grand Island, NY). Following an 18 hour incubation to allow for the maximal induction of PGHS-2, the conditioned medium was removed and assayed for PGE<sub>2</sub> activity by EIA analysis as described above.

10 U937 (ATCC, Rockville, MD) cells were grown in a similar fashion to the WISH cells. After incubation, the conditioned medium was removed and assayed for Cox-1 activity by EIA analysis as described above.

The data illustrating the inhibition of prostaglandin biosynthesis *in vitro* by compounds of this invention is shown in Table 2. U937 values indicate percent  
 15 inhibition at the particular micromolar dose level while parenthetical values indicate IC<sub>50</sub> values. Wish cell values indicate Cox-2 percent inhibition at the particular micromolar dose level.

**TABLE 2**

Example No.	U937 ( $\mu$ M)	Wish (mM)
7	(0.87)	(0.02)
10	62 @ 10	(0.11)
14	42 @ 1	34 @ 0.00001
15	33 @ 0.01	43 @ 0.001
17	41 @ 10	54 @ 1

20	60 @ 10	(0.13)
21	80 @ 10	(0.00004)
23	30 @ 10	62 @ 1

### Carrageenan Induced Paw Edema (CPE) in Rats

- Hindpaw edema was induced in male rats as described by Winter *et al.*,  
5 *Proc. Soc. Exp. Biol. Med.*, **1962**, *111*, 544. Briefly, male Sprague-Dawley rats  
weighing between 170 and 190 g were administered test compounds orally 1 hour  
prior to the subplantar injection of 0.1 ml of 1% sodium carrageenan (lambda  
carrageenan, Sigma Chemical Co., St Louis, MO) into the right hindpaw. Right  
paw volumes (ml) were measured immediately following injection of carrageenan  
10 for baseline volume measurements using a Buxco plethysmograph (Buxco  
Electronics, Inc., Troy, NY). Three hours after the injection of carrageenan, right  
paws were remeasured and paw edema calculated for each rat by subtracting the  
zero time reading from the 3 hour reading. Data are reported as mean percent  
inhibition +/- SEM. Statistical significance of results was analyzed by Dunnetts  
15 multiple comparison test where  $p < 0.05$  was considered statistically significant.

**Rat Carrageenan Pleural Inflammation Model (CIP)**

Pleural inflammation was induced in male adrenalectomized Sprague-Dawley rats following the method of Vinegar *et al.*, *Fed. Proc.* 1976, 35, 2447-2456. Animals were orally dosed with experimental compounds, in 0.2% HPMC, 30 minutes prior to the intrapleural injection of 2% lambda carrageenan (Sigma Chemical Co., St. Louis MO). Four hours later the animals were euthanized and the pleural cavities lavaged with ice cold saline. The lavage fluid was then added to two volumes of ice cold methanol (final methanol concentration 66%) to lyse cells and precipitate protein. Eicosanoids were determined by EIA as described above. The data illustrating the inhibition of prostaglandin biosynthesis *in vivo* by the compounds of this invention is shown in Table 3. Values reported are percent inhibition at 10 milligrams per kilogram body weight.

**TABLE 3**

Example No.	CIP Inh @10mpk
7	22
23	41

**15 Pharmaceutical Compositions**

The present invention also provides pharmaceutical compositions which comprise compounds of the present invention formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions of the present invention comprise a therapeutically effective amount of a compound of the present invention formulated together with one or more pharmaceutically acceptable carriers. As used herein, the term "pharmaceutically acceptable carrier" means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as

cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; 5 pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the 10 procedures and judgements well known to one skilled in the art. The pharmaceutical compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray.

15           The compounds of the present invention may be potentially useful in the treatment of several illness or disease states such as inflammatory diseases, dysmenorrhea, asthma, premature labor, osteoporosis, and ankylosing spondylitis. Current Drugs Ltd, ID Patent Fast Alert, AG16, May 9, 1997.

20           The compounds of the present invention may also be potentially useful in the treatment of cancers, and in particular, colon cancer. Proc. Natl. Acad. Sci., 94, pp. 3336-3340, 1997.

25           The compounds of the present invention may be useful by providing a pharmaceutical composition for inhibiting prostaglandin biosynthesis comprising a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt, ester, or prodrug thereof, and a pharmaceutically acceptable carrier.

30           The compounds of the present invention may be useful by providing a pharmaceutical composition for inhibiting prostaglandin biosynthesis comprising a therapeutically effective amount of a compound of formula II or a pharmaceutically acceptable salt, ester, or prodrug thereof; and a pharmaceutically acceptable carrier.

The compounds of the present invention may be useful by providing a pharmaceutical composition for inhibiting prostaglandin biosynthesis comprising a therapeutically effective amount of a compound of formula III or a pharmaceutically acceptable salt, ester, or prodrug thereof, and a pharmaceutically acceptable carrier.

The compounds of the present invention may be useful by providing a pharmaceutical composition for inhibiting prostaglandin biosynthesis comprising a therapeutically effective amount of a compound of formula IV or a pharmaceutically acceptable salt, ester, or prodrug thereof, and a pharmaceutically acceptable carrier.

In addition, the compounds of the present invention may be useful by providing a method for inhibiting prostaglandin biosynthesis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt, ester, or prodrug thereof.

The compounds of the present invention may be useful by providing a method for inhibiting prostaglandin biosynthesis comprising administering to a mammal in need of such treatment a therapeutically effective amount a compound of formula II or a pharmaceutically acceptable salt, ester, or prodrug thereof.

The compounds of the present invention may be useful by providing a method for inhibiting prostaglandin biosynthesis comprising administering to a mammal in need of such treatment a therapeutically effective amount compound of formula III or a pharmaceutically acceptable salt, ester, or prodrug thereof.

The compounds of the present invention may be useful by providing a method for inhibiting prostaglandin biosynthesis comprising administering to a mammal in need of such treatment a therapeutically effective amount a compound of formula IV or a pharmaceutically acceptable salt, ester, or prodrug thereof.

In addition, the compounds of the present invention may be useful by providing a method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula I.

In addition, the compounds of the present invention may be useful by providing a method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula II.

- 5 In addition, the compounds of the present invention may be useful by providing a method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula III.

- 10 In addition, the compounds of the present invention may be useful by providing a method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of formula IV.

- Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs.
- 15 In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (such as, for example, cottonseed,
- 20 groundnut, corn, germ, olive, castor, sesame oils, and the like), glycerol, tetrahydrofurfuryl alcohol, poly-ethyl-ene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

- 25 Injectable preparations, such as, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, such as, for
- 30 example, a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, isotonic sodium chloride solution, and the like. In addition, sterile, fixed oils are conventionally



employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectable preparations.

5 The injectable formulations can be sterilized by any method known in the art, such as, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

10 In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered  
15 drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsulated matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other  
20 biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention  
25 with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and thus melt in the rectum or vaginal cavity and release the active compound.

Solid dosage forms for oral administration include capsules, tablets, pills,  
30 powders, and granules. In such solid dosage forms, the active compound is usually mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as, for example, sodium citrate or dicalcium phosphate and/or a) fillers

or extenders such as, for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as, for example, glycerol, d) disintegrating agents such as, for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as, for example, paraffin, f) absorption accelerators such as, for example, quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as, for example, kaolin and bentonite clay, and) lubricants such as, for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients such as, for example, lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using excipients such as, for example, lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulation art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as, for example, sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as, for example, magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed

manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

5 Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

10 The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients such as, for example, animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

15 Powders and sprays can contain, in addition to the compounds of this invention, excipients such as, for example, lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons.

20 Transdermal patches have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in a suitable medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

25 According to the methods of treatment of the present invention, a patient, such as a human or mammal, is treated by administering to the patient a therapeutically effective amount of a compound of the invention, in such amounts and for such time as is necessary to achieve the desired result. By a "therapeutically effective amount" of a compound of the invention is meant a  
30 sufficient amount of the compound to provide the relief desired, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the

- present invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.
- 10 The total daily dose of the compounds of this invention administered to a human or other mammal in single or in divided doses can be in amounts, for example, from 0.001 to about 1000 mg/kg body weight daily or more preferably from about 0.1 to about 100 mg/kg body weight for oral administration or 0.01 to about 10 mg/kg for parenteral administration daily. Single dose compositions may contain such
- 15 amounts or submultiples thereof to make up the daily dose.


The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

- The reagents required for the synthesis of the compounds of the invention
- 20 are readily available from a number of commercial sources such as Aldrich Chemical Co. (Milwaukee, WI, USA); Sigma Chemical Co. (St. Louis, MO, USA); and Fluka Chemical Corp. (Ronkonkoma, NY, USA); Alfa Aesar (Ward Hill, MA 01835-9953); Eastman Chemical Company (Rochester, New York 14652-3512); Lancaster Synthesis Inc. (Windham, NH 03087-9977); Spectrum Chemical
- 25 Manufacturing Corp. (Janssen Chemical) (New Brunswick, NJ 08901); Pfaltz and Bauer (Waterbury, CT. 06708). Compounds which are not commercially available can be prepared by employing known methods from the chemical literature.

## CLAIMS

We claim:

- 5 1. A compound of formula I:



I

where

- X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic, heterocyclic (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;
- 15 R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy, alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl, haloalkynyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, aryl, arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy, arylcarbonylalkyl, heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy), heterocyclic (oxy), -C(O)R<sup>5</sup>, -C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -R<sup>6</sup>-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OH)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(OR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(NOR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C(NR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NOR<sup>d</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>CH(NR<sup>d</sup>R<sup>6</sup>)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX'<sub>3</sub>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(C X'<sub>2</sub>)<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, -(CH<sub>2</sub>)<sub>n</sub>(CHX')<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-CX'<sub>3</sub>, -(CH<sub>2</sub>)<sub>n</sub>(CHX')<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>,
- 20 25

wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, haloalkenyl, haloalkynyl, heterocyclic, and heterocyclic (alkyl);

wherein  $R^6$  is alkylene or alkenylene;

5  $R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl);


$R^{20}$  is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

10  $R^d$  and  $R^e$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl);

$X'$  is halogen;

$n$  is from 0 to about 10, and  $m$  is 0 to about 5;

15 at least one of  $R^1$ ,  $R^2$  and  $R^3$  is a group, substituted with a substituent having a group  $-X^1-R^9$ , having the formula:



where  $X^1$  is selected from the group consisting of  $-SO_2-$ ,  $-SO(NR^{10})-$ ,  $-PO(OR^{11})-$ , and  $-PO(NR^{12}R^{13})-$ ,

20  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino,  $-NHNH_2$ , alkylamino, dialkylamino, alkoxy, thiol, alkylthiol, and protecting groups,

$X^2$  is selected from the group consisting of hydrogen or halogen;

25  $R^{10}$ ,  $R^{11}$ ,  $R^{12}$ , and  $R^{13}$  are independently selected from the group consisting of hydrogen, alkyl, and cycloalkyl, or  $R^{12}$  and  $R^{13}$  can be taken together, with the nitrogen to which they are attached, to form a heterocyclic ring having from 3 to 6 atoms.


The remaining two of the groups of R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup>, are independently selected from the group consisting of hydrogen, hydroxy, hydroxyalkyl, halogen, alkyl, alkenyl, alkynyl, alkoxy, alkoxyalkyl, amido, amidoalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, amino, aminocarbonyl, aminocarbonylalkyl, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -C(R<sup>16</sup>)(R<sup>17</sup>)-,

C(O)NR<sup>21</sup>-, -C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl),

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; and


R<sup>21</sup> is selected from the group consisting of hydrogen, alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

2. A compound of the formula II:



II

wherein Z is a group having the formula:



where  $X^1$  is selected from the group consisting of  $-SO_2-$ ,  $-a\bar{n}$   
 $SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl,  
 alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino;

$X^2$  is selected from the group consisting of hydrogen or halogen;

5  $X$  is selected from the group consisting of O, S,  $NR^4$ ,  $N-OR^a$ , and  
 $N-NR^bR^c$ , wherein  $R^4$  is selected from the group consisting of alkyl, alkenyl,  
 cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic,  
 heterocyclic (alkyl), and arylalkyl; and  $R^a$ ,  $R^b$ , and  $R^c$  are independently selected  
 from the group consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;

10  $R$  is selected from the group consisting of hydrogen, alkyl, alkenyl,  
 alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy,  
 alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl,  
 haloalkynyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl,  
 cycloalkenylalkyl, aryl, arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy,  
 15 arylcarbonylalkyl, heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy),  
 heterocyclic (oxy),  $-C(O)R^5$ ,  $-(CH_2)_nC(O)R^5$ ,  $-(CH_2)_nC\equiv C-R^7$ ,  
 $-(CH_2)_n[CH(CX'_3)]_m-(CH_2)_n-CX'_3$ ,  $-(CH_2)_n(CX'_2)_m-(CH_2)_n-CX'_3$ ,  
 $-(CH_2)_n[CH(CX'_3)]_m-(CH_2)_n-R^8$ ,  $-(CH_2)_n(CX'_2)_m-(CH_2)_n-R^8$ ,  $-(CH_2)_n(CHX'_m)-$   
 $(CH_2)_n-CX'_3$ ,  $-(CH_2)_n(CHX'_m)-(CH_2)_n-R^8$ , and  $-(CH_2)_n-R^{20}$ ,

20 wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl,  
 alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and  
 heterocyclic (alkyl);

$R^7$  and  $R^8$  are independently selected from the group consisting of  
 hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl,  
 25 haloalkyl, heterocyclic, and heterocyclic (alkyl),

$R^{20}$  is selected from the group consisting of alkyl, alkenyl, haloalkyl,  
 cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

$X'$  is halogen;


$n$  is from 0 to about 10,  $m$  is from 0 to about 5;



R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of hydrogen, hydroxy, hydroxyalkyl, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -CH<sub>2</sub>-, -C(R<sup>16</sup>) (R<sup>17</sup>)-, -C(O)-, -C(O)O-, -NH-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and


R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

3. A compound according to claim 1 having the formula III:



wherein X, X<sup>1</sup>, X<sup>2</sup>, R, R<sup>1</sup>, R<sup>3</sup>, and R<sup>9</sup> are as defined above in Formula I; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

4. A compound of the formula III:



III

wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -  
 5 SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and  
 N-NR<sup>b</sup>RC, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cyclo-  
 10 alkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, aryl, heterocyclic, heterocyclic  
 (alkyl), and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group  
 consisting of alkyl, cycloalkyl, cycloalkylalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl,  
 alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl,  
 15 cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl,  
 arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, -(CH<sub>2</sub>)<sub>n</sub>C≡C-  
 R<sup>7</sup>, -(CH<sub>2</sub>)<sub>n</sub>[CH(CX<sup>3</sup>)]<sub>m</sub>-(CH<sub>2</sub>)<sub>n</sub>-R<sup>8</sup> and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl,  
 alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and  
 20 heterocyclic (alkyl);

R<sup>7</sup> and R<sup>8</sup> are independently selected from the group consisting of  
 hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl,  
 haloalkyl, heterocyclic, and heterocyclic (alkyl),

R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl,  
 25 cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

X<sup>3</sup> is halogen;

n is from 0 to about 10, m is from 0 to about 5;

R<sup>1</sup> and R<sup>3</sup> are independently selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -, -O-, -S-, -C(R<sup>16</sup>) (R<sup>17</sup>)-,

-C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

5. A compound according to claim 4 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen;

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, and -(CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

n is from 0 to about 10;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -, -O-, -S-, -C(R<sup>16</sup>)(R<sup>17</sup>)-,

-C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

6. A compound according to claim 4 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>RC<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), arylalkyl, and  $-(CH_2)_nC(O)R^5$ ;

5                    wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl); and

n is from 0 to about 10;

10                    $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and  $-Y-R^{14}$ , wherein Y is selected from the group consisting of,  $-O-$ ,  $-S-$ ,  $-C(R^{16})$  ( $R^{17}$ )-,

$-C(O)-$ ,  $-C(O)O-$ ,  $-NH-$ ,  $-NC(O)-$ , and  $-NR^{19}$ .  $R^{14}$  is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

15                    $R^{16}$ ,  $R^{17}$ , and  $R^{19}$  are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

20                   7. A compound according to claim 4 wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

$X^2$  is selected from the group consisting of hydrogen and halogen,

25                   X is selected from the group consisting of O, S,  $NR^4$ ,  $N-OR^a$ , and  $N-NR^bR^c$ , wherein  $R^4$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and  $R^a$ ,  $R^b$ , and  $R^c$  are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

- 5         $R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and  $-Y-R^{14}$ , wherein Y is selected from the group consisting of,  $-O-$ ,  $-S-$ ,  $-C(R^{16})(R^{17})-$ ,  $-C(O)-$ ,  $-C(O)O-$ ,  $-NH-$ ,  $-NC(O)-$ , and  $-NR^{19}-$ .  $R^{14}$  is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cyclo-
- 10    alkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

$R^{16}$ ,  $R^{17}$ , and  $R^{19}$  are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

15

8.        A compound according to claim 4 wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

- 20         $X^2$  is selected from the group consisting of hydrogen and halogen,

X is selected from the group consisting of O, S,  $NR^4$ ,  $N-OR^a$ , and  $N-NR^bR^c$ , wherein  $R^4$  is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and  $R^a$ ,  $R^b$ , and  $R^c$  are independently selected from the group consisting of alkyl, cyclo-

25    alkyl, alkylcycloalkyl, aryl, and arylalkyl;

R is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl); or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- 5 9. A compound according to claim 4 wherein X<sup>1</sup> is -SO<sub>2</sub>-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,


- 10 X is selected from the group consisting of O, S, NR<sup>4</sup>, N-OR<sup>a</sup>, and N-NR<sup>b</sup>R<sup>c</sup>, wherein R<sup>4</sup> is selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkylcycloalkyl, alkylcycloalkenyl, aryl, heteroaryl, and arylalkyl; and R<sup>a</sup>, R<sup>b</sup>, and R<sup>c</sup> are independently selected from the group consisting of alkyl, cycloalkyl, alkylcycloalkyl, aryl, and arylalkyl;

- 15 R is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and (CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup> where R<sup>20</sup> is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

- 20 R<sup>1</sup> and R<sup>2</sup> are independently selected from the group consisting of unsubstituted aryl and substituted aryl with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

10. A compounds of formula IV:



## IV

wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

5  $X^2$  is selected from the group consisting of hydrogen and halogen,

$R$  is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, alkoxy, alkoxyalkyl, carboxy, carboxyalkyl, cyanoalkyl, haloalkyl, haloalkenyl, haloalkynyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, cycloalkenyl, cycloalkenylalkyl, aryl, 10 arylalkyl, arylalkenyl, arylalkynyl, arylalkoxy, aryloxy, arylcarbonylalkyl, heterocyclic, heterocyclic (alkyl), heterocyclic (alkoxy), heterocyclic (oxy),  $-(CH_2)_nC(O)R^5$ ,  $-(CH_2)_nC\equiv C-R^7$ ,  $-(CH_2)_n[CH(CX'_3)]_m-(CH_2)_n-R^8$  and  $-(CH_2)_n-R^{20}$ ;

wherein  $R^5$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and 15 heterocyclic (alkyl);

$R^7$  and  $R^8$  are independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl),  $R^{20}$  is selected from the group 20 consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

$X'$  is halogen;

$n$  is from 0 to about 10,  $m$  is from 0 to about 5;

$R^1$  and  $R^2$  are independently selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, 25 amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl, heterocyclic, heterocyclic (alkyl), cyano, nitro, and  $-Y-R^{14}$ , wherein  $Y$  is selected from the group consisting of  $-, -O-, -S-, -C(R^{16})-(R^{17})-$ ,

$-C(O)-$ ,  $-C(O)O-$ ,  $-NH-$ ,  $-NC(O)-$ , and  $-NR^{19}-$ .  $R^{14}$  is selected from the group 30 consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and



R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

5

11. A compound according to claim 10 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

10 X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkenyl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>, and -  
15 (CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup>;

wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl);

20 R<sup>20</sup> is selected from the group consisting of alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkenyl, aryl, heterocyclic, and heterocyclic (alkyl);

n is from 0 to about 10;

R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, dialkylamino, arylamino, arylalkylamino, diarylamino, aryl,  
25 heterocyclic, heterocyclic (alkyl), cyano, nitro, and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -C(R<sup>16</sup>)(R<sup>17</sup>)-,

-C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

5

12. A compound according to claim 10 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

10 X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

R is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, alkylcarbonylalkyl, alkylsulfonylalkyl, alkylsulfonylarylalkyl, carboxyalkyl, cyanoalkyl, haloalkyl, hydroxyalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkynyl, heterocyclic, heterocyclic (alkyl), arylalkyl, and -(CH<sub>2</sub>)<sub>n</sub>C(O)R<sup>5</sup>;

15 wherein R<sup>5</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, arylalkyl, haloalkyl, heterocyclic, and heterocyclic (alkyl); and

n is from 0 to about 10;

20 R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and -Y-R<sup>14</sup>, wherein Y is selected from the group consisting of, -O-, -S-, -C(R<sup>16</sup>)(R<sup>17</sup>)-,

-C(O)-, -C(O)O-, -NH-, -NC(O)-, and -NR<sup>19</sup>-. R<sup>14</sup> is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cycloalkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

25 R<sup>16</sup>, R<sup>17</sup>, and R<sup>19</sup> are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

13. A compound according to claim 10 wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

5  $X^2$  is selected from the group consisting of hydrogen and halogen,

$R$  is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

$n$  is from 0 to about 10;

10  $R^1$  is hydrogen and  $R^2$  is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), and  $-Y-R^{14}$ , wherein  $Y$  is selected from the group consisting of,  $-O-$ ,  $-S-$ ,  $-C(R^{16})(R^{17})-$ ,

$-C(O)-$ ,  $-C(O)O-$ ,  $-NH-$ ,  $-NC(O)-$ , and  $-NR^{19}-$ .  $R^{14}$  is selected from the group consisting of hydrogen, halogen, alkyl, alkenyl, alkynyl, hydroxy, cycloalkyl, cyclo-

15 alkenyl, amino, cyano, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl), and

$R^{16}$ ,  $R^{17}$ , and  $R^{19}$  are independently selected from the group consisting of alkyl, alkenyl, cycloalkyl, cycloalkenyl, alkoxy, aryl, arylalkyl, heterocyclic, heterocyclic (alkyl), or cyano; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

20

14. A compound according to claim 10 wherein  $X^1$  is selected from the group consisting of  $-SO_2-$ , and  $-SO(NR^{10})-$ , and  $R^9$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

25  $X^2$  is selected from the group consisting of hydrogen and halogen,

$R$  is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and  $-(CH_2)_n-R^{20}$  where  $R^{20}$  is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

$n$  is from 0 to about 10;

R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of hydrogen, aryl, arylalkyl, heterocyclic, and heterocyclic (alkyl) ; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- 5           15. A compound according to claim 10 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>-, and -SO(NR<sup>10</sup>)-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

- 10           R is selected from haloalkyl, aryl, heterocyclic, heterocyclic (alkyl), and - (CH<sub>2</sub>)<sub>n</sub>-R<sup>20</sup> where R<sup>20</sup> is substituted and unsubstituted aryl wherein the substituted aryl compounds are substituted with halogen;

n is from 0 to about 10;

- 15           R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of hydrogen, aryl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine including, but not limited to, *p*-chlorophenyl, *p*-fluorophenyl, 3,4-dichlorophenyl, 3-chloro-4-fluoro-phenyl, and the like; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- 20           16. A compound according to claim 10 wherein X<sup>1</sup> is -SO<sub>2</sub>-, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

- 25           R is haloalkyl and R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of unsubstituted aryl and aryl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

- 30           17. A compound according to claim 10 wherein X<sup>1</sup> is selected from the group consisting of -SO<sub>2</sub>, and R<sup>9</sup> is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, amino, alkylamino, or dialkylamino;

X<sup>2</sup> is selected from the group consisting of hydrogen and halogen,

R is substituted and unsubstituted aryl and R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected from the group consisting of unsubstituted aryl and aryl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

18. A compound according to claim 10 wherein X<sup>1</sup> is SO<sub>2</sub>, R<sup>9</sup> is selected from alkyl and amino, X<sup>2</sup> is selected from the group consisting of hydrogen and halogen, R is substituted and unsubstituted aryl and R<sup>1</sup> is hydrogen and R<sup>2</sup> is selected unsubstituted aryl and aryl substituted with one, two, or three substituents selected from the group consisting of alkyl, alkoxy, fluorine and chlorine; or a pharmaceutically acceptable salt, ester, or prodrug thereof.

15

19. A compound according to claim 10, selected from the group consisting of:

- 5-(4-Methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-Benzyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 20 2-Methyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-Ethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-(4-Fluorobenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-(n-Butyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 25 2-(2,2,2-Trifluoroethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-(4-Fluoro- $\alpha$ -methylbenzyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-(n-Propyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 30 2-(n-Pentyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-Cyclohexylmethyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;
- 2-Phenacyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;

2-Propargyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-Cyclohexyl-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(2-Butynyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-pyridazinone;  
2-(Cyclobutanylmethyl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
5 pyridazinone; and  
2-(3-Methylbuten-2-yl)-5-(4-methylsulfonylphenyl)-6-(4-fluorophenyl)-3(2H)-  
pyridazinone; or a pharmaceutically acceptable salt or ester thereof.

10 20. A pharmaceutical composition for inhibiting prostaglandin  
biosynthesis comprising a therapeutically effective amount of the compound  
of claim 1 and a pharmaceutically acceptable carrier.

15 21. A pharmaceutical composition for inhibiting prostaglandin  
biosynthesis comprising a therapeutically effective amount of the compound  
of claim 2 and a pharmaceutically acceptable carrier.

20 22. A pharmaceutical composition for inhibiting prostaglandin  
biosynthesis comprising a therapeutically effective amount of the compound  
of claim 4 and a pharmaceutically acceptable carrier.

23. A pharmaceutical composition for inhibiting prostaglandin  
biosynthesis comprising a therapeutically effective amount of the compound  
of claim 10 and a pharmaceutically acceptable carrier.

25 24. A method for inhibiting prostaglandin biosynthesis comprising  
administering to a mammal in need of such treatment a therapeutically  
effective amount of a compound of claim 1.

25. A method for inhibiting prostaglandin biosynthesis comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claim 2.
- 5 26. A method for inhibiting prostaglandin biosynthesis comprising administering to a mammal a therapeutically effective amount of a compound of claim 4.
- 10 27. A method for inhibiting prostaglandin biosynthesis comprising administering to a mammal a therapeutically effective amount of a compound of claim 10.
- 15 28. A method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a therapeutically effective amount of a compound of claim 1.
- 20 29. A method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claim 1.
30. A method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal a therapeutically effective amount of a compound of claim 2.
- 25 31. A method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal a therapeutically effective amount of a compound of claim 4.

32. A method for treating pain, fever, inflammation, rheumatoid arthritis, osteoarthritis, and cancer comprising administering to a mammal a therapeutically effective amount of a compound of claim 10.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/17618

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6 C07D237/14 C07D409/04 C07F9/6509 C07D237/18 C07D237/20 A61K31/50				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D C07F A61K				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
A	EP 0 400 519 A (K.THOMAE) 5 December 1990  see page 1 - page 11; claims; examples 79-81  -----	1,10, 20-23		
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.				
* Special categories of cited documents : <table border="0"> <tr> <td style="vertical-align: top;">           "A" document defining the general state of the art which is not considered to be of particular relevance            "E" earlier document but published on or after the international filing date            "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)            "O" document referring to an oral disclosure, use, exhibition or other means            "P" document published prior to the international filing date but later than the priority date claimed         </td> <td style="vertical-align: top;">           "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention            "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone            "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.            "A" document member of the same patent family         </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family			
Date of the actual completion of the international search		Date of mailing of the international search report		
7 December 1998		17/12/1998		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Paternlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer		
		Francois, J		

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In ternational Application No

PCT/US 98/17618

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 400519 A	05-12-1990	DE 3917801 A	06-12-1990
		DE 3934436 A	18-04-1991
		AU 623262 B	07-05-1992
		AU 5612790 A	06-12-1990
		CA 2017957 A	01-12-1990
		DD 297969 A	30-01-1991
		JP 3236378 A	22-10-1991
		PT 94213 A	08-02-1991
		US 5135932 A	04-08-1992
-----			